



Salmonellosis

- Reported cases of disease, 1999-2003
- 5-year rate of disease, 1999-2003
- Reported cases by age group, 2003
- Reported cases by gender, 2003
- Reported cases by county, 2003
- Reported cases by month, 2003



Reported cases of Salmonellosis by age group, Rhode Island, 2003			
Age Group	Number of reported cases	Percent	Rate ¹ of disease per 100,000
< 1	12	8.8 %	98.3
1-4	19	13.9 %	36.8
5-9	17	12.4 %	23.6
10-19	11	8.0 %	7.5
20-29	16	11.7 %	11.7
30-39	17	12.4 %	10.6
40-49	23	16.8 %	14.3
50-59	10	7.3 %	8.6
60-74	11	8.0 %	9.8
75+	1	0.7 %	1.3
TOTAL	137	100.0%	13.1
¹ Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Salmonellosis by gender, Rhode Island, 2003			
Gender	Number of reported cases	Percent	Rate ¹ of disease per 100,000
Males	66	48.2%	13.1
Females	71	51.8 %	13.0
TOTAL	137	100.0%	13.1
¹ Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Salmonellosis by county, Rhode Island, 2003			
County	Number of reported cases	Percent	Rate ¹ of disease per 100,000
Bristol	4	2.9 %	7.9
Kent	27	19.7 %	16.2
Newport	13	9.5 %	15.2
Providence	68	49.6 %	10.9
Washington	16	11.7 %	7.3
Missing	9	6.6 %	----
TOTAL	137	100.0%	13.1
¹ Denominators for computing rates for RI were based on 2000 population counts obtained from the U.S. Census Bureau.			

**Reported cases of Salmonellosis ases,
by month, Rhode Island, 2003**

